

# Statistical Mechanical Ensembles and Typical Behavior of Macroscopic Systems

Joel L. Lebowitz


Departments of Mathematics and Physics  
Rutgers University

## Abstract

In this talk I will focus on describing, in a qualitative way, the reason why statistical mechanics is able to predict, with great certainty, behavior of macroscopic systems, both in equilibrium and out of it. I will relate this to the fact that this behavior is *typical* for systems represented by the usual Gibbs ensembles or those derived from them. These take small phase space volume to indicate small probability. I will not try to justify this here.


---

I thank S. Goldstein, E. Speer, and O. Penrose for very many, very helpful comments and A. Krueger for technical help with these slides. I also thank the NSF and the AFOSR for long term support.



The word typical is used in ordinary language to denote a state or behavior usually observed in a given situation. Example: glasses at room temperature are typically hard. To make this more precise we need to specify a probability distribution  $P$  over some space  $\Omega$  of elements  $X$ . We can then say that an event such as  $f(X) = C$  is typical if it has probability close to 1 wrt  $P$ .


When  $f = C$  is typical and  $P$  is "uniform" over  $\Omega$ , I shall say that " $Y$  is typical of  $\Omega$ ", meaning  $f(Y) = C$ .



For macroscopic systems an overwhelming majority of the microstates in the micro-canonical ensemble give the same “thermodynamic” behavior: such behavior is typical.


In fact, the fraction of systems with noticeable macroscopic deviations from the average behavior, computed in such an ensemble, is exponentially small in the number of degrees of freedom of the system.

The functions on the phase space which correspond to such thermodynamic behavior will be described later.



Analogous statements hold for the (exponentially small) subsets of the micro-canonical ensemble which describe systems in nonequilibrium macrostates (to be defined below), wrt to the microcanonical measure restricted to such subsets.

In particular, this applies to subsets for which the typical behavior is the time asymmetric approach to equilibrium, encoded in the second law and observed in individual macroscopic systems.



Once one accepts the applicability of these measures (ensembles) to physical systems they can be used to predict the typical behavior of individual macroscopic systems *and* not just some average behavior.

The observed behavior does not require explanations based on ergodicity, time averaging, or subjective information theory.

I will begin with classical systems where statistical mechanics was first developed by Boltzmann, Maxwell and their contemporaries. In fact the macroscopic behavior I will consider here is pretty much the same for classical as for quantum systems.

## Classical Systems

In classical mechanics, the microstate of a system of  $N$  particles confined to a region  $V$  in  $\mathbb{R}^d$  is a point  $X$  in the  $2dN$ -dimensional phase space,  $\Gamma$ ,

$$X = (\mathbf{r}_1, \mathbf{v}_1, \dots, \mathbf{r}_N, \mathbf{v}_N), \quad \mathbf{r}_i \in V \subset \mathbb{R}^d, \quad \mathbf{v}_i \in \mathbb{R}^d \quad (1)$$

Its time evolution is given by a Hamiltonian  $H(X)$  which conserves energy, so  $X(t) = T_t X$  will be confined to the energy surface  $H(X) = E$ , to which we will always refer by  $\Gamma_E$ , a thin shell surrounding that surface.

A macroscopic system is one with “very large”  $N$ , say  $N \gtrsim 10^{20}$ .

## Time Evolution

The kind of Hamiltonian I have in mind is the usual effective one for most atoms,

$$H(X) = \frac{1}{2} \sum_{i=1}^N \mathbf{v}_i^2 + \sum_{i<j} \phi(r_{ij}) \quad (2)$$

with  $r_{ij} = |\mathbf{r}_i - \mathbf{r}_j|$  and  $\phi(r)$  a potential of the Lennard-Jones or hard sphere type. The time evolution of the microstate is given by the solution of the equations

$$\frac{d\mathbf{r}_i}{dt} = \mathbf{v}_i, \quad \frac{d\mathbf{v}_i}{dt} = - \sum_{j \neq i} \frac{\partial \phi(r_{ij})}{\partial \mathbf{r}_i}, \quad (3)$$

This time evolution,  $T_t X = X(t)$ , is reversible: given  $X = (\mathbf{r}_1, \mathbf{v}_1, \dots, \mathbf{r}_N, \mathbf{v}_N)$  define  $RX = (\mathbf{r}_1, -\mathbf{v}_1, \dots, \mathbf{r}_N, -\mathbf{v}_N)$  then

$$T_s R T_s X = RX. \quad (4)$$

## Macrostates


To describe the macroscopic state of such a system of  $N$  particles in a box  $V$ , we make use of a much cruder description than that provided by the microstate  $X$ .

We shall denote by  $M$  such a macroscopic description, and by  $M(X)$  the macrostate of the system in the microstate  $X$ .

The exact definition of macrostates is somewhat dependent on how precise a macroscopic description we are interested in. In all cases the state  $M$  of macroscopic systems will depend only on functions which are sums of terms which depend only on the coordinates and momenta of a few particles. I shall refer to these as thermodynamic functions.

The amount of information in  $M(X)$  is much smaller than that contained in the microstate  $X$ .





As an example we may divide  $V$  into  $K$  cells, where  $K$  is large but still  $K \ll N$ , and define  $M$  by specifying the number of particles, the momentum and the amount of energy in each cell, with some tolerance. Clearly there are many  $X$ 's (in fact a continuum) which correspond to the same  $M$ .

Let  $\Gamma_M$  be the region in  $\Gamma_E$  consisting of all microstates  $X$  corresponding to a given macrostate  $M$  and denote by  $|\Gamma_M|$  its phase-space volume.

Taking  $H(X)$  of the form (2) it can be shown<sup>1</sup> that in every  $\Gamma_E$  of a macroscopic system there is one region  $\Gamma_M$  which has most of the volume of  $\Gamma_E$ . This is called the equilibrium macrostate  $M_{\text{eq}}$ ,

$$\frac{|\Gamma_{M_{\text{eq}}}|}{|\Gamma_E|} = 1 - \varepsilon \quad (5)$$

with  $\varepsilon \ll 1$ . A system is then in macroscopic thermal equilibrium iff  $X \in \Gamma_{M_{\text{eq}}}$ . When  $M(X)$  specifies a nonequilibrium state,  $|\Gamma_M|$  is very much smaller.

Thus for a gas consisting of  $N$  particles in a volume  $V$  the ratio of  $|\Gamma_M|$ , the volume of a macrostate  $M$  in which all the particles are in the left half of the box and  $|\Gamma_{M_{\text{eq}}}|$ , the volume of the macrostate  $M_{\text{eq}}$  in which there are  $(\frac{1}{2} \pm 10^{-6})N$  particles in the left half of the box, is of order  $2^{-N}$ ; see Figure 1.

---

<sup>1</sup>La73





Figure 1: Schematic Picture of the decomposition of  $\Gamma_E$ . Here  $\Gamma_{eq} \equiv \Gamma_{M_{eq}}$ .

The second picture is slightly more faithful.

Neither shows the topology or differences in relative sizes of the different  $\Gamma_M$ 's. In general, the closer  $M$  is to  $M_{eq}$  the larger  $\Gamma_M$ : e.g. 60% vs 100% of particles in the left half of the box.



The thermodynamic functions have approximately the same value for almost all  $X \in \Gamma_{M_{\text{eq}}}$ , hence they are typical for  $\Gamma_{M_{\text{eq}}}$  and observed for (almost) all systems in equilibrium.

In fact since  $|\Gamma_{M_{\text{eq}}}| \simeq |\Gamma_E|$  they are also typical of  $X \in \Gamma_E$ .

This justifies the use of the microcanonical ensemble for  $N \gg 1$  to compute relevant properties of an equilibrium system, independent of whether or not the dynamics is ergodic in a mathematical sense.

## Nonequilibrium States

Thus, most microstates in  $\Gamma_E$  of a macroscopic system correspond to the system being in equilibrium.

As we shall see when we discuss quantum systems below, a similar statement is true for most wave functions, in fact also for most energy eigenstates, in  $\mathcal{H}_E$ , the energy shell of the Hilbert space.

Fortunately there are also microstates which correspond to macroscopic systems which are out of equilibrium (or we would not be here).


Consider now the time evolution of the macrostate  $M(X(t))$  when the system is in the microstate  $X(t_0) \in \Gamma_M$ ,  $M \neq M_{\text{eq}}$ , at an “initial” time  $t_0$ , and is isolated for  $t > t_0$ .

## Approach to Equilibrium

Boltzmann (also Maxwell, Kelvin, ...) argued that given the disparity in the sizes of the  $\Gamma_M$  corresponding to the various macrostates, the evolution of a microstate  $X(t_0)$ , typical wrt a uniform measure on  $\Gamma_M$ , will be such that  $|\Gamma_{M(X(t))}|$  will not decrease (on a macroscopic scale) for  $t > t_0$ .

Thus the evolution towards equilibrium of macroscopic systems which start in the macrostate  $\Gamma_M$ ,  $M \neq M_{\text{eq}}$ , and are kept (effectively) isolated afterwards, is typical with respect to the micro-canonical measure restricted to  $\Gamma_M$ .

N.B. We have assumed that such measures give the right probabilities for physical systems.



In fact one can argue that for any  $\Gamma_M$  the relative volume of the set of microstates  $X \in \Gamma_M$  for which the above is false over some macroscopic time period,  $\tau$ , goes to zero exponentially in the number of atoms in the system.  $\tau$  should be long enough for the macrostate to undergo a macroscopically noticeable change but not longer than the age of the universe.

## Boltzmann's Entropy

To make a connection with the Second Law, Boltzmann defined the (Boltzmann) entropy of a macroscopic system in a microstate  $X$  as

$$S_B(X) = \log |\Gamma_{M(X)}| = S_B(M(X)). \quad (6)$$


The entropy of the equilibrium macrostate

$$S_B(M_{\text{eq}}) = \log |\Gamma_{M_{\text{eq}}}| \simeq \log |\Gamma_E| \quad (7)$$

is then the maximum value of  $S_B(M)$  over all macrostates  $M$  with energy  $E$ .

Boltzmann then showed that  $S_B(M_{\text{eq}})$  agrees with the Clausius thermodynamic entropy of a gas in equilibrium.





Boltzmann's heuristic argument for the non-decrease of entropy, based on relative phase space volume, is the correct explanation for the behavior typically observed in actual macroscopic systems. It is, however, very far from a mathematical theorem and contains no quantitative information about time scales<sup>2</sup>.

A proof would be provided by the rigorous derivation from the microscopic dynamics of the kinetic and hydrodynamic equations commonly used to describe the time asymmetric, entropy increasing behavior of macroscopic systems out of equilibrium.

This has been achieved so far only for the Boltzmann equation for dilute gases. This was done rigorously (in appropriate limits) by Oscar Lanford in 1975.


---

<sup>2</sup>As Mark Kac said: an argument convinces a reasonable person, a proof convinces a stubborn one

## The Boltzmann Equation for Dilute Gases

Following Boltzmann, we refine the description of a macrostate  $M$  by noting that the microstate  $X = \{\mathbf{r}_i, \mathbf{v}_i\}$ ,  $i = 1, \dots, N$ , can be considered as a set of  $N$  points in six dimensional one particle space.


Dividing up this one particle space into cells  $\Delta_\alpha$ , centered on  $(\mathbf{r}_\alpha, \mathbf{v}_\alpha)$ , of volume  $|\Delta_\alpha|$ , we can describe the macro (meso) states of the system  $X \in \Gamma_E$  by specifying, with some leeway, the fraction of particles in each  $\Delta_\alpha$ .



Conversely given a smooth distribution  $f(\mathbf{x}, \mathbf{v})$  in the one-particle space we define the macrostate  $M_f$  and the region  $\Gamma_{M_f} \subset \Gamma_E$  as consisting of phase points  $X$  such that the fraction of particles in each  $\Delta_\alpha$ , is given by

$$N_\alpha(X)/N \cong \int_{\Delta_\alpha} d\mathbf{x}d\mathbf{v} f(\mathbf{x}, \mathbf{v}). \quad (8)$$

Since we are dealing with a dilute gas we can neglect the interaction energy so  $f$  determines the energy  $E$  of the macrostate  $M_f$ .




Boltzmann then used his deep physical intuition (and hints from Maxwell) to derive his eponymous equation for the time evolution of the macrostate  $M_f$ .

The reasoning behind Boltzmann's analysis and the exact derivation of the Boltzmann equation (BE) is carefully explained in Lanford's beautiful non-technical article<sup>3</sup>. This is one of the best mathematical-physics articles I have ever read. I strongly recommend it.

I will now give a bird's eye view of what I think is the essence of that article.

---

<sup>3</sup>La76



Consider a gas consisting of  $N$  spheres of diameter  $d$  in a volume  $V \subset \mathbb{R}^3$  evolving according to Hamiltonian dynamics with elastic collisions.

Keeping  $V$  fixed consider now a sequence of systems with different particle numbers, and different diameters  $d$ , such that  $N \rightarrow \infty$ ,  $d \rightarrow 0$ , while  $Nd^2 \rightarrow b > 0$  (and so  $Nd^3 \rightarrow 0$ ). This is called the Boltzmann-Grad (BG) limit.

Let  $f_0(\mathbf{x}, \mathbf{v}) \geq 0, \mathbf{x} \in V \subset \mathbb{R}^3, \mathbf{v} \in \mathbb{R}^3$ , be a smooth distribution of integral one in the one particle space.

Consider now for each  $N$  all the phase points  $X_N$  of this gas such that  $X_N \in \Gamma_{M_{f_0}} \subset \Gamma_E$ . The dimension of  $\Gamma_E$  (and  $\Gamma_{M_{f_0}}$ ) will be  $6N$  but  $f_0(\mathbf{x}, \mathbf{v})$  remains fixed. As  $N$  increases the fraction of particles in  $\Delta_\alpha$ , is required satisfy ever more closely  $N_\alpha(X_N)/N \cong \int_{\Delta_\alpha} f_0(\mathbf{x}, \mathbf{v}) d\mathbf{x}d\mathbf{v}$ , with equality in the BG limit,

$$\lim_{\text{BG}} N_\alpha(X_N)/N = \int_{\Delta_\alpha} f_0(\mathbf{x}, \mathbf{v}) d\mathbf{x}d\mathbf{v} \quad (9)$$

Lanford's theorem then says:

Given  $f_0$ , a typical microstate  $X_N(0) \in \Gamma_{M_{f_0}}$  will evolve, via the Hamiltonian dynamics, to a microstate  $X_N(t)$  such that for all reasonable sets of  $\Delta_\alpha$ 's

$$\lim_{BG} \frac{N_\alpha(X_N(t))}{N} = \int_{\Delta_\alpha} f(\mathbf{x}, \mathbf{v}, t) d\mathbf{x} d\mathbf{v} \quad (10)$$

where  $f(\mathbf{x}, \mathbf{v}, t)$  solves the Boltzmann equation

$$\frac{\partial f}{\partial t} + \mathbf{v} \cdot \frac{\partial f}{\partial \mathbf{x}} = Q(f, f) \quad (11)$$

with initial condition  $f_0(\mathbf{x}, \mathbf{v})$ .

Here again typical is with respect to the uniform measure on  $\Gamma_{M_{f_0}}$ .

The BE gives a deterministic evolution from  $M_{f_0}$  to  $M_{f_t}$  in the BG limit. It will hold to a good approximation for a macroscopic dilute gas. It accurately describes the typical time evolution of the “empirical” distribution of atoms in the one particle space of such an individual system.

The Boltzmann entropy of a dilute gas in a macrostate  $M_f$  (as defined in (6)) is,

$$S_B(f) = S_B(M_f) = \log |\Gamma_{M_f}| \quad (12)$$

where  $|\Gamma_{M_f}|$  is for an  $N$  particle system the  $6N$  dimensional phase space volume of  $\Gamma_{M_f}$

$S_B(f)$  was actually computed by Boltzmann. He showed that, up to constants, this is given for a dilute gas of  $N$  particles by

$$\frac{1}{N} S_B(f) = - \int_V d\mathbf{x} \int_{\mathbb{R}^3} d\mathbf{v} f(\mathbf{x}, \mathbf{v}) \log f(\mathbf{x}, \mathbf{v}). \quad (13)$$

While (13) looks like, and is sometimes mistakenly confused with, the Gibbs-Shannon entropy per particle for a system in a product measure of the form  $\prod_i f(\mathbf{x}_i, \mathbf{v}_i)$ , it is conceptually not the same at all.  $S_B(f)$  is the entropy of an individual system with empirical distribution  $f(\mathbf{x}, \mathbf{v})$  (see below).



The maximum of  $S_B(f)$  over all  $f$  for a system of  $N$  particles in a unit volume with a given energy, which is for a dilute gas just the kinetic energy, is given by the Maxwell distribution

$$f_{\text{eq}} = (2\pi kT/m)^{-3/2} \exp[-m\mathbf{v}^2/2kT] \quad (14)$$

where  $kT = 2/3(E/N)$ .

The distribution  $f_{\text{eq}}(\mathbf{x}, \mathbf{v})$  is the unique stationary solution of the BE.

In this case  $\Gamma_{M_{f_{\text{eq}}}} = \Gamma_{M_{\text{eq}}}$  and the entropy/particle is given by

$$\frac{1}{N} S_B(f_{\text{eq}}) = \frac{3}{2} \log T - \log \frac{N}{|V|} + \text{Const.} \quad (15)$$

the same as the equilibrium Clausius entropy for a dilute gas, with density  $\frac{N}{|V|}$ .

The second law now says that for a *typical* microstate of a dilute gas with  $X(0) \in \Gamma_{M_{f_0}}$ ,  $f_0 \neq f_{\text{eq}}$ ,  $f(\mathbf{x}, \mathbf{v}, t) = f_t$  has to be such that  $S_B(f_t) \geq S_B(f_{t'})$ , for  $t \geq t'$ .

This is exactly what happens for  $f(\mathbf{x}, \mathbf{v}, t)$  evolving according to the Boltzmann equation:


$$\frac{d}{dt} S_B(f_t) \geq 0, \quad \text{Boltzmann's } \mathcal{H}\text{-theorem} \quad (16)$$

As put by Boltzmann<sup>4</sup>:

*“In one respect we have even generalized the entropy principle here, in that we have been able to define the entropy in a gas that is not in a stationary state.”*

---

<sup>4</sup>Bo98



Lanford could prove his theorem only for a time  $0 < t < \tau$ . This is still the case at present. In fact one can only prove the existence of smooth solutions of the BE by using Lanford's theorem and then of course only for times  $t < \tau$ .

The time  $\tau$  for which Lanford's theorem holds is about one fifth of the mean free time between collisions, but that is a purely technical problem. This time is long enough for the Boltzmann entropy per particle to increase by a finite amount.

## More General Macroscopic Equations

Going beyond the BE suppose, more generally, that the time evolution of the macrostate  $M$ , given by  $M(X(t)) = M_t$ , effectively satisfies an autonomous deterministic time asymmetric equation, such as the diffusion or the heat equation. (I shall consider here for simplicity macrostates  $M$  which are invariant under velocity reversal.)

Having such an equation like the Boltzmann equation just discussed means that if  $t_3 > t_2 > t_1$ , then the microscopic dynamics  $T_t$  carries  $\Gamma_{M_{t_1}}$  inside  $\Gamma_{M_{t_2}}$  and  $\Gamma_{M_2}$  inside  $\Gamma_3$ , i.e.  $T_{t_2-t_1}\Gamma_{M_{t_1}} \subset \Gamma_{M_{t_2}}$  and  $T_{t_3-t_2}\Gamma_{M_{t_2}} \subset \Gamma_{M_{t_3}}$ , with *negligible error*. Put otherwise a typical phase point in  $\Gamma_{M_{t_1}} = \Gamma_{M_1}$  will go to  $\Gamma_{M_2}$  and then to  $\Gamma_{M_3}$ , i.e.  $T_{t_3-t_1}\Gamma_{M_1} \subset \Gamma_{M_3}$ .





Figure 2: Time evolution of  $\Gamma_{M_1}$




The fact that phase space volume is conserved by the Hamiltonian time evolution implies that  $|\Gamma_{M_1}| \leq |\Gamma_{M_2}| \leq |\Gamma_{M_3}|$ , and thus that  $S_B(M_3) \geq S_B(M_2) \geq S_B(M_1)$ .

Hence the solution of any deterministic macroscopic equation for  $M_t$  has to satisfy the inequality<sup>5</sup>  $\frac{d}{dt} S_B(M_t) \geq 0$ , exactly what Boltzmann showed for the BE.


---

<sup>5</sup>Go04,Pe70



We note that the existence of such a macroscopic equation implies (at least insofar as the macro-variables are concerned) that the phase points in the region in  $\Gamma_2$  coming from  $\Gamma_1$  behave, forward in time, as microstates typical of  $\Gamma_2$ .

They are, however, very atypical backwards in time: if we reverse all the velocities in  $\Gamma_2$ , then at a later time,  $t' = t_2 + (t_2 - t_1)$  all of the points initially in  $\Gamma_{M_1}$  will again be in  $\Gamma_{M_1}$  (with their velocities reversed) a smaller region than  $\Gamma_{M_2}$ . We would thus have a violation of the second law.



The reason for this asymmetry in typical behaviors is due to initial conditions. That is, when nature or the experimentalist who is part of nature, starts out at  $t_1$  with a nonequilibrium system in an initial state  $X \in \Gamma_{M_1}$  we can assume, for  $X$  typical of  $\Gamma_{M_1}$ , that  $X(t)$  will continue to be typical of  $M_t$  insofar as times greater than  $t_1$  are concerned but obviously not with regard to times earlier than  $t_1$ .

This was shown explicitly in the Lanford derivation of the BE: reversing the velocities at some  $t > 0$  ( $t < \tau$ ) violates the assumptions on the initial conditions required for the derivation of the Boltzmann equation.



But what about real life situations?

What corresponds to an appropriate choice of initial time and initial conditions, one than can somehow make plausible that at that time the state  $X$  is atypical (because it is a low entropy nonequilibrium state)? Somewhat surprisingly, if one thinks hard about it, one is pushed to consider the very beginning of the universe we live in.

This would correspond according to our current physical theories to the time just after the “Big Bang”. The importance of initial conditions, Big Bang or not, was already fully understood by Boltzmann and others as the quotes below show.




## Initial Conditions

*“From the fact that the differential equations of mechanics are left unchanged by reversing the sign of time without changing anything else, Herr Ostwald concludes that the mechanical view of the world cannot explain why natural processes always run preferentially in a definite direction. **But such a view appears to me to overlook that mechanical events are determined not only by differential equations, but also by initial conditions. In direct contrast to Herr Ostwald I have called it one of the most brilliant confirmations of the mechanical view of Nature that it provides an extraordinarily good picture of the dissipation of energy, as long as one assumes that the world began in an initial state satisfying certain conditions. I have called this state an improbable state.**”*

— L. Boltzmann<sup>6</sup>

---

<sup>6</sup>Bo97




*“It is necessary to add to the physical laws the hypothesis that, in the past the universe was more ordered in the technical sense, [i.e. low  $S_B$ ] than it is today ... to make an understanding of irreversibility.”*

— R.P. Feynman<sup>7</sup>


---

<sup>7</sup>Fe67



**Figure 3:** “Creation of the universe: a fanciful description! The Creator’s pin has to find a tiny box, just 1 part in  $10^{10^{123}}$  of the entire phase-space volume, in order to create a universe with as special a Big Bang as we actually find.” from R. Penrose, *The Emperor’s New Mind*

The “tiny box” in the figure is a macrostate with low  $S_B$ . N.B. It is not necessary to select a particular microstate. Almost all microstates in a low-entropy macrostate will behave in a similar way.



It may be relevant to mention here a question I was asked during a talk I gave on the subject:

Q: What does the initial state of the universe have to do with the fact that when I put my sugar cube in my tea it dissolves irreversibly?

A: Nothing directly. But the fact that you, the sugar cube and the tea are all here is a consequence of the initial low entropy state of the universe.

## Boltzmann vs. Gibbs Entropies

Given an ensemble (probability) density  $\mu(X)$ ,  $X \in \Gamma$ , the Gibbs-Shannon entropy is given by

$$S_\mu \equiv - \int_\Gamma \mu \log \mu \, dX. \quad (17)$$

Clearly if  $\mu = \tilde{\mu}_M$ , where

$$\tilde{\mu}_M = \begin{cases} |\Gamma_M|^{-1}, & \text{if } X \in \Gamma_M; \\ 0, & \text{otherwise} \end{cases} \quad (18)$$

then

$$S_G(\tilde{\mu}_M) = \log |\Gamma_M| = S_B(M). \quad (19)$$

This is essentially the case for the microcanonical ensemble since  $\Gamma_E \simeq \Gamma_{M_{\text{eq}}}$ . By the equivalence of ensembles for macroscopic systems the same is true for the canonical and other Gibbs ensembles.

Thus the Gibbs and Boltzmann entropies are equal for equilibrium systems.

However, as  $\mu = \mu_t$  evolves via the Hamiltonian dynamics for isolated systems  $S_G(\mu)$  does not change in time.  $S_G(\mu)$  is therefore “useless” for such systems not in equilibrium, while  $S_B(M(X_t))$  captures the essence of typical macroscopic behavior. In particular it satisfies the second law of thermodynamics.



**Figure 4:** Boltzmann's grave in Zentralfriedhof, Vienna, with bust and entropy formula

## Summary of Boltzmann's Ideas (also Maxwell, Kelvin, ...)

Time-asymmetric behavior as embodied in the second law of thermodynamics is observed in *individual macroscopic* systems. It can be understood as arising naturally from time-symmetric microscopic laws when account is taken of a) the great disparity between microscopic and macroscopic sizes, b) initial conditions, and c) that what we observe is “typical” behaviors — not all imaginable ones. Common alternate explanations, such as those based on equating irreversible macroscopic behavior with ergodic or mixing properties of ensembles (probability distributions) already present for chaotic dynamical systems having only a few degrees of freedom or on the impossibility of having a truly isolated system, are either unnecessary, misguided or misleading.

Let me end this section on systems described by classical mechanics by quoting Einstein's tribute to Boltzmann.

*“On the basis of kinetic theory of gases Boltzmann had discovered that, aside from a constant factor, entropy is equivalent to the logarithm of the “probability” of the state under consideration. Through this insight he recognized the nature of course of events which, in the sense of thermodynamics, are “irreversible”. Seen from the molecular-mechanical point of view, however all courses of events are reversible. If one calls a molecular-theoretically defined state a microscopically described one, or, more briefly, micro-state, then an immensely large number ( $Z$ ) of states belong to a macroscopic condition.  $Z$  is then a measure of the probability of a chosen macro-state. **This idea appears to be of outstanding importance also because of the fact that its usefulness is not limited to microscopic description on the basis of mechanics.”***

— A. Einstein, Autobiographical notes




## Quantum Systems

Let me turn now to quantum systems. These are very different on the microscopic level from, but quite similar on the macroscopic level to, classical systems.


The first question is: what is the best quantum description of the microstate of a physically isolated macroscopic system? Unfortunately there is no consensus on this even a century after the birth of quantum mechanics and its perfect success in explaining and predicting microscopic phenomena.

There is however no doubt that, as put by Einstein, Boltzmann's ideas are applicable to the quantum (real) world.



I will assume for this talk that the microstate of a macroscopic system with an energy in a narrow interval  $(E, E + \Delta E)$  to be a wave function  $\psi \in \mathcal{H}_E$ , where  $\mathcal{H}_E$  is a shell in Hilbert space of thickness  $\Delta E$ , i.e.  $\mathcal{H}_E$  consists of all linear combination of energy eigenfunctions in the range  $(E, E + \Delta E)$ ,  $\Delta E \ll E$  but very large compared to the spacing between levels.

This is not perfect (problems with Schrödinger's Cat) but it will have to do for the present (see below). Using a density matrix with eigenstates in  $\mathcal{H}_E$  would give similar results.



The next question then is which  $\psi$  correspond to the system being in macroscopic thermal equilibrium (MATE), i.e. what is the analog of a classical microstate  $X$  lying in  $\Gamma_{M_{\text{eq}}}$ .

Following von Neumann<sup>8</sup>, we take the macro-observables corresponding to the macrostate  $M$  to commute with each other. We assume that this can be achieved by suitably “rounding off”, i.e. coarse-graining, the operators representing the macro-observables.

---

<sup>8</sup>Chapter 5 of Ne55

The coarse-grained energy operator commutes with the other coarse-grained macro-observables, which I shall also denote by  $M$ . Thus all  $M$ 's can be regarded as operators on  $\mathcal{H}_E$ . Their joint spectral decomposition defines an orthogonal decomposition

$$\mathcal{H}_E = \bigoplus_{\nu} \mathcal{H}_{\nu}, \quad (20)$$

Each subspace  $\mathcal{H}_{\nu}$  consists of all linear combinations of the joint eigenvectors of the macro-observables whose eigenvalues lie in a narrow range. These subspaces  $\mathcal{H}_{\nu}$  correspond to the different macrostates and the orthogonal decomposition (20) of  $\mathcal{H}_E$  corresponds to the division of the classical energy shell  $\Gamma_E$  into disjoint regions  $\Gamma_M$ .



A system is in a macrostate  $M_\nu$  if its wave function  $\psi$  is “close” to  $\mathcal{H}_\nu$ , i.e.

$$\langle \psi | P_\nu | \psi \rangle \geq 1 - \delta, \quad \delta \ll 1, \quad (21)$$

with  $P_\nu$  being the projection to  $\mathcal{H}_\nu$ .

As noted earlier, due to the Schrödinger’s Cat problem, there will be  $\psi$ ’s which are a superposition of  $\psi$ ’s in different macrostates.

To remedy this one has to go beyond the Copenhagen doctrine that the wave function is a complete description<sup>9</sup>. For the present let me say that I would interpret such  $\psi$ ’s when they arise physically as giving probabilities  $\langle \psi | P_\nu | \psi \rangle$  of being in different macrostates,  $M_\nu$ .

---

<sup>9</sup>Be87,Go98

The analog of the phase space volume  $|\Gamma_M|$  is the dimension  $d_\nu$  of each macro space  $\mathcal{H}_\nu$ .


As in the classical case, it is generally true that one of the  $\mathcal{H}_\nu$ , denoted  $\mathcal{H}_{\text{eq}}$ , has most of the dimensions of  $\mathcal{H}_E$ , i.e.,

$$\frac{\dim \mathcal{H}_{\text{eq}}}{\dim \mathcal{H}_E} = 1 - \varepsilon \quad (22)$$

with  $\varepsilon \ll 1$ .

The Boltzmann entropy  $S_B$  of a system in a macrostate  $M_\nu$  is given by the log of the dimension of the macro space  $\mathcal{H}_\nu$ ;

$$S_B(M_\nu) = \log d_\nu. \quad (23)$$



Given this correspondence between partitioning  $\Gamma_E$  and decomposing  $\mathcal{H}_E$  most everything we said about isolated classical macroscopic systems then also holds for quantum systems. In particular for  $\varepsilon \ll \delta$ , most  $\psi \in \mathcal{H}_E$  are in MATE.


The Boltzmann argument for increase in entropy of isolated macroscopic systems out of equilibrium is then similar to that in the classical case<sup>10</sup>.

When I say “most  $\psi$ ”, I mean with respect to a uniform measure on the unit sphere in  $\mathcal{H}_E$ . This measure on “wave functions” was considered already by Schrödinger and particularly by Felix Bloch<sup>11</sup>. It yields the microcanonical density matrix  $\rho^{\text{mc}}$  but goes beyond it, in ascribing probabilities to any subset of  $\mathcal{H}_E$ .

---

<sup>10</sup>Gr94

<sup>11</sup>Go06



There is however also something beyond MATE which can be used for the characterization of quantum systems in equilibrium.

Unlike classical systems, where any subsystem of a system in a microstate  $X$  is also in a microstate  $X_S$ , having definite positions and velocities, a subsystem  $S$  of a quantum system with a wave function  $\psi$  will usually not be described by a wave function, but rather by a density matrix  $\rho_S^\psi$ , where

$$\rho_S^\psi = \text{tr}_{S^c} |\psi\rangle\langle\psi| \quad (24)$$

is the reduced density matrix of  $S$  obtained by tracing out the complement  $S^c$  of  $S$ .



We can now define<sup>12</sup> a system with wave function  $\psi$  to be in microscopic thermal equilibrium (MITE) if, for any not-too-large subsystem  $S$ , say subsystems with linear dimension  $\ell < \ell_0$ , for an appropriate  $\ell_0$ , the reduced density matrix of  $S$  is close to the thermal equilibrium density matrix of  $S$

$$\rho_S^\psi \approx \rho_S^{\text{mc}}, \quad (25)$$

$$\rho_S^{\text{mc}} = \text{tr}_{S^c} \rho^{\text{mc}}, \quad (26)$$

where  $\rho^{\text{mc}}$  is the microcanonical density matrix corresponding to a uniform distribution over energy eigenstates in  $\mathcal{H}_E$ . For macroscopic systems  $\rho_S^{\text{mc}}$  can be replaced by  $\rho_S^{\text{ca}}$ , where  $\rho^{\text{ca}}$  is the canonical density matrix.

The distinction between MITE and MATE is important for systems with many-body localization (MBL) for which most, if not all, the energy eigenfunctions fail to be in MITE while necessarily most of them, but not all, are in MATE.

---

<sup>12</sup>Go17

The argument for most energy eigenfunctions being in MATE is based on the fact that, calling  $D$  the dimension of  $\mathcal{H}_E$ , we have for energy eigenfunctions  $|n\rangle$

$$\frac{1}{D} \sum_{n=1}^D \langle n | P_{\nu_{\text{eq}}} | n \rangle = \frac{1}{D} \text{tr}(P_{\nu_{\text{eq}}}) = 1 - \varepsilon \quad (27)$$

Noting that  $\langle n | P_{\nu_{\text{eq}}} | n \rangle \leq 1$ , the average being close to 1 means that most eigenstates are close to  $\mathcal{H}_{\nu_{\text{eq}}}$ . This is consistent with the Eigenstate Thermalization Hypothesis (ETH).

In fact for generic macroscopic systems, including those with MBL, most wave functions in an energy shell are in both MATE and MITE.

This follows from the following result.

## Canonical Typicality

Consider an isolated system consisting of two parts. Call them system 1 and 2 or system and reservoir. Then<sup>13</sup> we have the following result.

Let  $H$  be the Hamiltonian of the whole system and let the number of particles in system 1 and 2 be  $N_1 \ll N_2$ . Let  $\mathcal{H}_E \subset \mathcal{H}_1 \otimes \mathcal{H}_2$  be an energy shell. Then for most  $\psi \in \mathcal{H}_E$  with  $\|\psi\| = 1$ ,


$$\text{tr}_2 |\psi\rangle\langle\psi| \approx \text{tr}_2 \rho^{\text{mc}}, \quad (28)$$

where  $\rho^{\text{mc}}$  is the microcanonical density matrix of the whole system at energy  $E$ , i.e. equal weight to all energy eigenstates in  $\mathcal{H}_E$ .

When the interaction between systems 1 and 2 is weak then, as is well known,  $\text{tr}_2 \rho^{\text{mc}} \approx \frac{1}{Z} e^{-\beta H_1}$  for  $\beta = \beta(E) = dS_{\text{eq}}(E)/dE$ .

---


<sup>13</sup>Go06,Le08



The theorem says that most wave functions in the energy shell  $\mathcal{H}_E$  are both in MATE and in MITE. In fact for macroscopic systems one can show that MITE implies MATE.

The opposite is however not true.

This is particularly relevant when one considers energy eigenfunctions  $|n\rangle$ . While most energy eigenstates, including those for systems with MBL must, as shown, generally be in MATE, most energy eigenfunctions for systems with MBL are not in MITE.



Interestingly MITE has been found experimentally and computationally to hold for systems with Hilbert spaces with dimension as small as 100 or so, for which the notion of macrostates is not really appropriate.

This brings up the question of how to apply what I have discussed here to systems with a small number of degrees of freedom. When considering such systems it is relevant to remember that probabilities of  $\mathcal{O}(10^{-6})$  might not be so different in practice from probabilities of  $\mathcal{O}(10^{-20})$ .

I will not discuss this or the related issues of “stochastic thermodynamics” here.

If MATE-ETH holds strictly, i.e., if *all* energy eigenstates in  $\mathcal{H}_E$  are in MATE, then every state  $\psi \in \mathcal{H}_E$  will sooner or later reach MATE and spend most of the time in MATE in the long run. That is because, writing  $\overline{f(t)} = \lim_{T \rightarrow \infty} \frac{1}{T} \int_0^T f(t) dt$  for time averages,  $|n\rangle$  for the energy eigenstate with eigenvalue  $E_n$  and  $\psi_t = e^{-iHt}\psi$ ,

$$\overline{\langle \psi_t | P_{\text{eq}} | \psi_t \rangle} = \sum_{n, n'} \overline{\langle \psi | n \rangle e^{iE_n t} \langle n | P_{\text{eq}} | n' \rangle e^{-iE_{n'} t} \langle n' | \psi \rangle} \quad (29)$$

$$= \sum_n |\langle \psi | n \rangle|^2 \langle n | P_{\text{eq}} | n \rangle \geq \sum_n |\langle \psi | n \rangle|^2 (1 - \delta) \quad (30)$$


$$= 1 - \delta, \quad (31)$$

provided  $H$  is non-degenerate, i.e.,  $E_n \neq E_{n'}$  for  $n \neq n'$  (using  $\overline{e^{iEt}} = 1$  if  $E = 0$  and  $= 0$  otherwise).

A similar statement is true when there is degeneracy.

## References

- [Be87] J.S. Bell (1987). *Speakable and Unspeakable in Quantum Mechanics*, Cambridge University Press, Cambridge, England.
- [Bo97] L. Boltzmann (1897). On Zermelo's Paper "On the Mechanical Explanation of Irreversible Processes". *Annalen der Physik*, **60**, 392-398.
- [Bo98] L. Boltzmann (1898). *Vorlesungen über Gastheorie*. Leipzig: Barth (Part I, Part II 1898). English translation by S.G. Brush: *Lectures on Gas Theory*. Berkeley: University of California Press (1964).
- [Ce94] C. Cercignani, R. Illner, M. Pulvirenti (1994). *The Mathematical Theory of Dilute Gases*. Berlin: Springer-Verlag.



[Fe67] R.P. Feynman (1967). *The Character of Physical Law*. Chapter 5. MIT Press, Cambridge, Mass.

[Go04] S. Goldstein, J.L. Lebowitz (2004). On the (Boltzmann) entropy of nonequilibrium systems. *Physica D*, **193**, 53-66.

[Go06] S. Goldstein, J.L. Lebowitz, R. Tumulka, N. Zanghi (2006). Canonical typicality. *Physical Review Letters*, **96**, 050403.

[Go98] S. Goldstein (1998). Quantum theory without observers. *Physics Today*, Part One: March, 42-46. Part Two: April, 38-42.




[Go17] S. Goldstein, D.A. Huse, J.L. Lebowitz, R. Tumulka (2017). Macroscopic and microscopic thermal equilibrium. *Annalen der Physik*, **529**, 1600301.

[Gr94] R. Griffiths (1994). Statistical Irreversibility: Classical and Quantum. Pages 147-159 in J.J. Halliwell, J. Pérez-Mercader, and W.H. Zurek (editors): *Physical Origin of Time Asymmetry*. Cambridge University Press.

[La73] O.E. Lanford (1973). Entropy and Equilibrium States in Classical Mechanics. Pages 1-113 in A. Lenard (editor): *Statistical Mechanics and Mathematical Problems*, Lecture Notes in Physics vol. **2**. Berlin:Springer-Verlag.

[La76] O.E. Lanford (1976). On a derivation of the Boltzmann equation. *Astérisque*, **40**, 117-137. Reprinted in J.L. Lebowitz and E.W. Montroll: *Nonequilibrium Phenomena — The Boltzmann Equation*, North-Holland (1983).



[Le08] J.L. Lebowitz (2008). From Time-symmetric Microscopic Dynamics to Time-asymmetric Macroscopic Behavior: An Overview. Pages 63-88 in G. Gallavotti, W.L. Reiter, J. Yngvason (editors): *Boltzmann's Legacy*. Zürich: European Mathematical Society.

[Ne55] J. von Neumann (1955). *Mathematical Foundations of Quantum Mechanics*. Pages 398-416. Princeton University Press, Princeton. Translated from German ed. (1932: Springer, Berlin) by R.T. Beyer.

[Pe70] O. Penrose (1970). *Foundations of Statistical Mechanics*. Pergamon, Elmsford, N.Y. Reprinted by Dover (2005).

[Pe90] R. Penrose (1990). *The Emperor's New Mind*. Chapter 7. Oxford University Press.